

P8269**Ablative vs. nonablative LASER technologies in aging hands: A randomized trial for a new treatment perspective**

Valeria Campos, MD, MS, Faculdade de Medicina de Jundiá (FMJ), Jundiá, Brazil; Fabiola Schauffler, MD, Faculdade de Medicina de Jundiá (FMJ), Jundiá, Brazil; Luciana Maluf, MD, Faculdade de Medicina de Jundiá (FMJ), Jundiá, Brazil; Luz Grohs, MD, Faculdade de Medicina de Jundiá (FMJ), Jundiá, Brazil; Marcella Martins, MD, Faculdade de Medicina de Jundiá (FMJ), Jundiá, Brazil; Mariana Wanczinski, MD, Faculdade de Medicina de Jundiá (FMJ), Jundiá, Brazil; Paulo Cunha, MD, PhD, Faculdade de Medicina de Jundiá (FMJ), Jundiá, Brazil

Background: Aged hands is a condition typically characterized by wrinkles, dermal atrophy, skin laxity, solar lentigines, actinic keratoses, seborrheic keratoses, and prominent veins. The search for effective treatments for rejuvenation hands has led to the use of LASER technologies and IPL systems as well.

Objective: To assess the efficacy of 2940-nm and 1340-nm LASER devices and IPL system for overall improvement of skin quality, collagen regeneration and also depigmentation on the dorsal surfaces of the hands.

Methods: Twelve patients with brown macules and sun damage spots on dorsal hands were treated. Both dorsal hands were split in 2 equal areas each. Lateral-side part of the right hand was used for controlling and was not treated, while 2940-nm LASER device was performed over medium side of the same hand. On left dorsal hand, 1340-nm LASER device was used on the lateral-side part, while on the medium side both 1340nm LASER and IPL system combined treatment was performed. Patients were treated twice with 4-week intervals. Treatment photos and informed consent were obtained. Pre- and posttreatment pictures, clinical data assessment investigation, and also patient questionnaires were collected for data analysis.

Results: Good to excellent results for rejuvenating hands were observed with all light-source systems. Remodeling of collagen and neocollagen and normalizing the mottled hyperpigmentation were mostly observed on the medium surface of the left hand with 1340-nm LASER plus IPL sources 92% (11/12 patients). The majority of patients, about 83% (10/12), do prefer this treatment modality while 2 of them do prefer the 2940-nm LASER treatment. Most common reported side effects were mild erythema, longer-lasting edema, blister, and pruritus.

Conclusion: Although all of the light devices had been effective for skin rejuvenation of the dorsal hands, 1340-nm LASER plus IPL system were the best treatment option in this study not only for overall efficacy, by stimulating collagen but also improving the discromia, but also for its safety assessment. Additional studies are also required to confirm these findings.

Commercial support: None identified.

P8585**Efficacy and patient satisfaction of a new nonablative fractional Nd:YAP laser 1340 nm for facial rejuvenation in Brazilian patients**

Valeria Campos, MD, MS, Faculdade de Medicina de Jundiá (FMJ), Jundiá, Brazil; Juliana Jordão, MD, Hospital Universitário Evangélico de Curitiba, Curitiba, Brazil

This present study wants to demonstrate a new fractional nonablative Nd:YAP 1340 nm laser technology that has been safely and effectively applied to treat photodamage in different skin types. Efforts to attend a growing demand of patients who seeks a procedure that treats wrinkles and improves skin textures with minimal to none downtime has been led to a constant upgrade in nonablative fractional lasers technologies. Intend of study is to determine the safety, efficacy, and patient satisfaction of a new nonablative fractional Nd:YAP laser 1340 nm in facial skin rejuvenation of Brazilian skin types. A comparison was made between treatments with low fluence and multiple passes versus treatments with higher fluence in a single pass. Twenty subjects with visible cutaneous photodamage, all with skin phototypes II to VI, were treated with a new fractional nonablative laser handpiece (Etherea 1340 ProDeep; INDUSTRA Technologies, São Carlos, Brazil). All patients returned every 4 weeks for a total of 3 sessions and were followed up with monthly photos and grading until 6 months after their last treatment session. They also were asked to fill out a "severity scale" where levels of pigmentation, rhytides, skin tone and tightness, texture, and patient satisfaction were noted after last treatment session. Pigmentation, rhytides, skin tone and tightness, and texture were also evaluated by 2 physicians not involved in the study and by Canfield VISIA Complexion. All patients answered a questionnaire at the end of evaluation. Postevaluation questionnaire showed that 100% of the patients felt improvement and 75% would recommend the same treatment to a friend. A new fractional nonablative Nd:YAP 1340 nm laser can be safely and effectively performed to treat photodamage in darker skin types. Although most patients preferred the single pass laser treatment, we did not observe significant difference in the outcome after a low fluence multiple pass versus high fluence single pass treatment and the incidence of side effects were higher in the high fluence single pass treatment. Additional studies with a larger number of subjects and biopsy specimens for histologic assessment are required.

Commercial support: None identified.

P7999**Efficacy of 1064 nm Q-switched Nd:YAG laser in the treatment of nevus of Ota in patients with Fitzpatrick skin types IV and V and utility of melanin index in the treatment outcome**

Sethuraman Gomathy, MD, All India Institute of Medical Sciences, New Delhi, India; Sreenivas V., All India Institute of Medical Sciences, New Delhi, India; Vinod K. Sharma, MD, All India Institute of Medical Sciences, New Delhi, India

Background: Q-switched Nd:YAG laser has been used in the treatment of nevus of Ota in all skin types with variable success rate. Data with an objective assessment parameter to this laser treatment is lacking.

Objective: To determine the efficacy and safety of the Q-switched Nd:YAG laser (1064-nm) in the treatment of nevus of Ota in Fitzpatrick skin type IV and V and also to evaluate the utility of melanin index in assessing the treatment response.

Methods: Patients treated with 1064 nm Nd:YAG laser at an interval of 6 to 10 weeks, for a period of 1 to 2 years were studied. The objective improvement (pigment clearance) was determined by melanin index from 2 fixed points: A1, 2 cm below the pupil at the midpupillary line (when the gaze is fixed); A2, the most prominent part of zygoma. The melanin index in these 2 areas was recorded as M1 and M2 respectively. The subjective clinical improvement was determined by the physician and the patient global assessment score.

Results: Thirty-five patients of nevus of Ota were included in the study. All excepting 1 patient had unilateral involvement. The mean baseline melanin indices M1 and M2 were 59.54 ± 9.72 and 59.02 ± 9.16 , respectively. At the last visit the mean M1 and M2 decreased to 53.8 ± 8.55 ($P < .001$) and 54.13 ± 6.01 ($P < .001$), respectively. Patient global assessment showed that overall 26 (74.3%) had >50% pigment clearance from the baseline. Fifteen patients (42.8%) had 50% to 75% improvement, 10 patients (28.6%) had 75% to 95% improvement and only 1 (2.9%) patient had >95% improvement. Assessment performed by the dermatologist blinded to the laser sessions and parameters, showed that 20 patients (57.1%) had >50% improvement. Eight patients (22.9%) had 50% to 75% improvement, 11 patients (31.4%) had 75% to 95% improvement and only 1 (2.9%) patient had >95% improvement. Spotty hypopigmentation after laser treatment was seen in 8 (22.9%) patients which improved over a period of time in most of them. Hyperpigmentation seen in 2 patients also reduced gradually.

Conclusion: The 1064 nm Q-switched Nd:YAG laser offers good improvement in patients with nevus of Ota in darker skin types IV/V. The melanin index, a simple noninvasive parameter is useful in assessing the treatment response more objectively.

Commercial support: None identified.

P8575**Fractional ablative LASER versus fractional ablative and coagulative LASER for treatment of photodamaged skin in arms and forearms**

Valeria Campos, MD, MS, Faculdade de Medicina de Jundiá (FMJ), Jundiá, Brazil; Clovis Pinto, MD, Faculdade de Medicina de Jundiá (FMJ), Jundiá, Brazil; Flavia Trevisan, MD, Faculdade de Medicina de Jundiá (FMJ), Jundiá, Brazil; Juliana Jordão, MD, Hospital Universitário Evangélico de Curitiba, Curitiba, Brazil; Luiza Pitassi, MD, MS, UNICAMP, Campinas, Brazil; Talita Santos, MD, Faculdade de Medicina de Jundiá (FMJ), Jundiá, Brazil

Photodamaged skin in arms and forearms are a common issue in dermatologic offices. Treatment with topical creams, peeling, and nonablative fractional lasers have mild results. Ablative lasers showed to have achieved the best results. The main issue, however, relies on extra facial areas, by increasing the risk of adverse effects caused by ablative lasers enhancement. The aim of this present study is to compare the whole effects of a fractional pure-ablative laser versus a fractional ablative and coagulative laser to treat photodamaged skin in arms and forearms, by using low energy doses in order to prevent side effects. Twenty patients have received a pure-ablative fractional laser treatment (known as single mode laser) in 1 arm and forearm, and in the other arm and forearm a combination of both ablative and coagulative laser treatment (known as dual mode laser). Both sides have been treated with a single pass by using a 2940 nm Er:YAG fractional laser (ETHEREA 2940 DualMode, INDUSTRA Technologies, Brazil), receiving 2 treatment sessions with 1 month of between sessions. In single mode was used 500-us pulse width with 10 mJ/MTZ fluence, and 500 us/3 ms pulse width with 10 mJ/MTZ and 10 J/cm² fluences for tx in dual mode. Patients were submitted to a skin biopsy before treatment, 1 month after the first and the second treatment, and 3 months after second treatment. Clinical pictures were recorded pretreatment and 3 months of posttreatment. Fifty percent of the patients have preferred the results of dual-mode treatment sessions. The pain was related to be more intense in this side in all of patients, as well. All patients noted a clear improvement in both sides. Biopsy has shown an epidermal and dermal collagen thickening after both treatments, but a more prominent result were observed in dual mode treated areas. No side effects were noticed in any treatment options. The use of low energy of fractional pure-ablative and fractional ablative and coagulative lasers are a safe and effective treatment option to treat photodamaged arms and forearms.

Commercial support: None identified.